6.6


What you should learn

GOAL Find the rational zeros of a polynomial function.

GOAL 2 Use polynomial equations to solve real-life problems, such as finding the dimensions of a monument in Ex. 60.

Why you should learn it

▼ To model real-life quantities, such as the volume of a representation of the Louvre pyramid in



Finding Rational Zeros

GOAL

USING THE RATIONAL ZERO THEOREM

The polynomial function

$$f(x) = 64x^3 + 120x^2 - 34x - 105$$

has $-\frac{3}{2}$, $-\frac{5}{4}$, and $\frac{7}{8}$ as its zeros. Notice that the numerators of these zeros (-3, -5, and 7) are factors of the constant term, -105. Also notice that the denominators (2, 4, 4)and 8) are factors of the leading coefficient, 64. These observations are generalized by the rational zero theorem.

THE RATIONAL ZERO THEOREM

If $f(x) = a_n x^n + \cdots + a_1 x + a_0$ has integer coefficients, then every rational zero of f has the following form:

$$\frac{p}{q} = \frac{\text{factor of constant term } a_0}{\text{factor of leading coefficient } a_n}$$

EXAMPLE 1

Using the Rational Zero Theorem

Find the rational zeros of $f(x) = x^3 + 2x^2 - 11x - 12$.

SOLUTION

List the possible rational zeros. The leading coefficient is 1 and the constant term is -12. So, the possible rational zeros are:

$$x = \pm \frac{1}{1}, \pm \frac{2}{1}, \pm \frac{3}{1}, \pm \frac{4}{1}, \pm \frac{6}{1}, \pm \frac{12}{1}$$

Test these zeros using synthetic division.

Since -1 is a zero of f, you can write the following:

$$f(x) = (x+1)(x^2 + x - 12)$$

Factor the trinomial and use the factor theorem.

$$f(x) = (x + 1)(x^2 + x - 12) = (x + 1)(x - 3)(x + 4)$$

The zeros of f are -1, 3, and -4.

In Example 1, the leading coefficient is 1. When the leading coefficient is not 1, the list of possible rational zeros can increase dramatically. In such cases the search can be shortened by sketching the function's graph—either by hand or by using a graphing calculator.

EXAMPLE 2 Using the Rational Zero Theorem


Find all real zeros of $f(x) = 10x^4 - 3x^3 - 29x^2 + 5x + 12$.

SOLUTION

List the possible rational zeros of $f: \pm \frac{1}{1}, \pm \frac{2}{1}, \pm \frac{3}{1}, \pm \frac{4}{1}, \pm \frac{6}{1}, \pm \frac{12}{1}, \pm \frac{3}{2}, \pm \frac{1}{5}, \pm \frac{2}{5}, \pm \frac{3}{5}, \pm \frac{6}{5}, \pm \frac{12}{5}, \pm \frac{1}{10}, \pm \frac{3}{10}, \pm \frac{12}{10}.$

Choose values to check

With so many possibilities, it is worth your time to sketch the graph of the function. From the graph, it appears that some reasonable choices are $x = -\frac{3}{2}$, $x = -\frac{3}{5}$, $x = \frac{4}{5}$, and $x = \frac{3}{2}$.



Check the chosen values using synthetic division.

Factor out a binomial using the result of the synthetic division.

$$f(x) = \left(x + \frac{3}{2}\right)(10x^3 - 18x^2 - 2x + 8)$$
 Rewrite as a product of two factors.
$$= \left(x + \frac{3}{2}\right)(2)(5x^3 - 9x^2 - x + 4)$$
 Factor 2 out of the second factor.
$$= (2x + 3)(5x^3 - 9x^2 - x + 4)$$
 Multiply the first factor by 2.

Repeat the steps above for $g(x) = 5x^3 - 9x^2 - x + 4$.

Any zero of g will also be a zero of f. The possible *rational* zeros of g are $x = \pm 1, \pm 2, \pm 4, \pm \frac{1}{5}, \pm \frac{2}{5}$, and $\pm \frac{4}{5}$. The graph of f shows that $\frac{4}{5}$ may be a zero.

So
$$f(x) = (2x + 3)\left(x - \frac{4}{5}\right)(5x^2 - 5x - 5) = (2x + 3)(5x - 4)(x^2 - x - 1).$$

Find the remaining zeros of f by using the quadratic formula to solve $x^2 - x - 1 = 0$.

The real zeros of f are
$$-\frac{3}{2}, \frac{4}{5}, \frac{1+\sqrt{5}}{2}$$
, and $\frac{1-\sqrt{5}}{2}$.


GOAL 2 SOLVING POLYNOMIAL EQUATIONS IN REAL LIFE



EXAMPLE 3

Writing and Using a Polynomial Model


You are designing a candle-making kit. Each kit will contain 25 cubic inches of candle wax and a mold for making a model of the pyramidshaped building at the Louvre Museum in Paris, France. You want the height of the candle to be 2 inches less than the length of each side of the candle's square base. What should the dimensions of your candle mold be?



SOLUTION

The volume is $V = \frac{1}{3}Bh$ where B is the area of the base and h is the height.





Volume = 25LABELS Side of square base = x

> Area of base = x^2 (square inches)

Height = x - 2

(inches)

(inches)

ALGEBRAIC MODEL

x-values.

$$25 = \frac{1}{3} x^2 (x - 2)$$
 Write algebraic model.


(cubic inches)


$$75 = x^3 - 2x^2$$

Multiply each side by 3 and simplify.

$$0 = x^3 - 2x^2 - 75$$

Subtract 75 from each side.





Use the possible solutions. Note that in this case, it makes sense to test only positive

← 5 is a solution.

So x = 5 is a solution. The other two solutions, which satisfy $x^2 + 3x + 15 = 0$, are $x = \frac{-3 \pm i\sqrt{51}}{2}$ and can be discarded because they are imaginary.

The base of the candle mold should be 5 inches by 5 inches. The height of the mold should be 5 - 2 = 3 inches.





I.M. PEI designed the pyramid at the Louvre. His geometric architecture can be seen in Boston, New York, Dallas, Los Angeles, Taiwan, Beijing, and Singapore.

GUIDED PRACTICE

Vocabulary Check ✓

Concept Check

1. Complete this statement of the rational zero theorem: If a polynomial function has integer coefficients, then every rational zero of the function has the form $\frac{p}{q}$, where p is a factor of the $\underline{?}$ and q is a factor of the $\underline{?}$.

2. For each polynomial function, decide whether you can use the rational zero theorem to find its zeros. Explain why or why not.

a.
$$f(x) = 6x^2 - 8x + 4$$
 b. $f(x) = 0.3x^2 + 2x + 4.5$ **c.** $f(x) = \frac{1}{4}x^2 - x + \frac{7}{8}$

3. Describe a method you can use to shorten the list of possible rational zeros when using the rational zero theorem.

Skill Check

List the possible rational zeros of f using the rational zero theorem.

4.
$$f(x) = x^3 + 14x^2 + 41x - 56$$
 5. $f(x) = x^3 - 17x^2 + 54x + 72$

6.
$$f(x) = 2x^3 + 7x^2 - 7x + 30$$
 7. $f(x) = 5x^4 + 12x^3 - 16x^2 + 10$

Find all the real zeros of the function.

8.
$$f(x) = x^3 - 3x^2 - 6x + 8$$
 9. $f(x) = x^3 + 4x^2 - x - 4$

10.
$$f(x) = 2x^3 - 5x^2 - 2x + 5$$
 11. $f(x) = 2x^3 - x^2 - 15x + 18$

12.
$$f(x) = x^3 + 4x^2 + x - 6$$
 13. $f(x) = x^3 + 5x^2 - x - 5$

14. Suppose you have 18 cubic inches of wax and you want to make a candle in the shape of a pyramid with a square base. If you want the height of the candle to be 3 inches greater than the length of each side of the base, what should the dimensions of the candle be?

PRACTICE AND APPLICATIONS

STUDENT HELP

Extra Practice to help you master skills is on p. 948.

STUDENT HELP

► HOMEWORK HELP

Example 1: Exs. 15–32 **Example 2**: Exs. 33–58

Example 3: Exs. 59-64

LISTING RATIONAL ZEROS List the possible rational zeros of f using the rational zero theorem.

15.
$$f(x) = x^4 + 2x^2 - 24$$

17.
$$f(x) = 2x^5 + x^2 + 16$$

19.
$$f(x) = 6x^4 - 3x^3 + x + 10$$

21.
$$f(x) = 8x^2 - 12x - 3$$

16.
$$f(x) = 2x^3 + 5x^2 - 6x - 1$$

18.
$$f(x) = 2x^3 + 9x^2 - 53x - 60$$

20.
$$f(x) = 4x^3 + 5x^2 - 3$$

22.
$$f(x) = 3x^4 + 2x^3 - x + 15$$

USING SYNTHETIC DIVISION Use synthetic division to decide which of the following are zeros of the function: 1, -1, 2, -2.

23.
$$f(x) = x^3 + 7x^2 - 4x - 28$$

25.
$$f(x) = x^4 + 3x^3 - 7x^2 - 27x - 18$$

27.
$$f(x) = x^4 + 3x^3 + 3x^2 - 3x - 4$$

29.
$$f(x) = x^3 - 3x^2 + 4x - 12$$

31.
$$f(x) = x^6 - 2x^4 - 11x^2 + 12$$

24.
$$f(x) = x^3 + 5x^2 + 2x - 8$$

26.
$$f(x) = 2x^4 - 9x^3 + 8x^2 + 9x - 10$$

28.
$$f(x) = 3x^4 + 3x^3 + 2x^2 + 5x - 10$$

30.
$$f(x) = x^3 + x^2 - 11x + 10$$

32.
$$f(x) = x^5 - x^4 - 2x^3 - x^2 + x + 2$$

FINDING REAL ZEROS Find all the real zeros of the function.

33.
$$f(x) = x^3 - 8x^2 - 23x + 30$$

33.
$$f(x) = x^3 - 8x^2 - 23x + 30$$
 34. $f(x) = x^3 + 2x^2 - 11x - 12$ **35.** $f(x) = x^3 - 7x^2 + 2x + 40$ **36.** $f(x) = x^3 + x^2 - 2x - 2$

37.
$$f(x) = x^3 + 72 - 5x^2 - 18x$$

38.
$$f(x) = x^3 + 9x^2 - 4x - 36$$

39.
$$f(x) = x^4 - 5x^3 + 7x^2 + 3x - 10$$

40.
$$f(x) = x^4 + x^3 + x^2 - 9x - 10$$

41.
$$f(x) = x^4 + x^3 - 11x^2 - 9x + 18$$

42.
$$f(x) = x^4 - 3x^3 + 6x^2 - 2x - 12$$

41.
$$f(x) = x^{5} + x^{4} - 9x^{3} - 5x^{2} - 36$$


42.
$$f(x) = x^4 - 3x^3 + 6x^2 - 2x - 12$$


44. $f(x) = x^5 - x^4 - 7x^3 + 11x^2 - 8x + 12$

ELIMINATING POSSIBLE ZEROS Use the graph to shorten the list of possible rational zeros. Then find all the real zeros of the function.

45.
$$f(x) = 4x^3 - 12x^2 - x + 15$$

46.
$$f(x) = -3x^3 + 20x^2 - 36x + 16$$





FINDING REAL ZEROS Find all the real zeros of the function.

47.
$$f(x) = 2x^3 + 4x^2 - 2x - 4$$

48.
$$f(x) = 2x^3 - 5x^2 - 14x + 8$$

49.
$$f(x) = 2x^3 - 5x^2 - x + 6$$

50.
$$f(x) = 2x^3 + x^2 - 50x - 25$$

51.
$$f(x) = 2x^3 - x^2 - 32x + 16$$

52.
$$f(x) = 3x^3 + 12x^2 + 3x - 18$$

53.
$$f(x) = 2x^4 + 3x^3 - 3x^2 + 3x - 5$$

54.
$$f(x) = 3x^4 - 8x^3 - 5x^2 + 16x - 5$$

55.
$$f(x) = 2x^4 + x^3 - x^2 - x - 1$$

56.
$$f(x) = 3x^4 + 11x^3 + 11x^2 + x - 2$$


x ft

$$\mathbf{E7} \ f(x) = 2x^5 + x^4 + 22x + 16$$

EQ
$$f(x) = 2x^5 + x^4 + 242x = 91$$

57.
$$f(x) = 2x^5 + x^4 - 32x - 16$$


58.
$$f(x) = 3x^5 + x^4 - 243x - 81$$



$$S = 10t^3 + 115t^2 + 25t + 2505$$

where *S* is the sales (in millions of dollars) and *t* is the number of years since 1990. In what year were about \$3885 million of health products sold? (Hint: First substitute 3885 for S, then divide both sides by 5.)

- **60. MONUMENTS** You are designing a monument and a base as shown at the right. You will use 90 cubic feet of concrete for both pieces. Find the value of x.
- **61.** MOLTEN GLASS At a factory, molten glass is poured into molds to make paperweights. Each mold is a rectangular prism whose height is 3 inches greater than the length of each side of the square base. A machine pours 20 cubic inches of liquid glass into each mold. What are the dimensions of the mold?



Ex. 60

solving in Ex. 60.

STUDENT HELP

HOMEWORK HELP Visit our Web site www.mcdougallittell.com for help with problem





FOCUS ON APPLICATIONS




SAND SCULPTURE
The tallest sand
sculptures built were over
20 feet tall and each
consisted of hundreds
of tons of sand.

- **62.** SAND CASTLES You are designing a kit for making sand castles. You want one of the molds to be a cone that will hold 48π cubic inches of sand. What should the dimensions of the cone be if you want the height to be 5 inches more than the radius of the base?
- 63. SWIMMING POOLS You are designing an in-ground lap swimming pool with a volume of 2000 cubic feet. The width of the pool should be 5 feet more than the depth, and the length should be 35 feet more than the depth. What should the dimensions of the pool be?





64. WHEELCHAIR RAMPS You are building a solid concrete wheelchair ramp. The width of the ramp is three times the height, and the length is 5 feet more than 10 times the height. If 150 cubic feet of concrete is used, what are the dimensions of the ramp?





QUANTITATIVE COMPARISON In Exercises 65 and 66, choose the statement that is true about the given quantities.

- (A) The quantity in column A is greater.
- **B** The quantity in column B is greater.
- **©** The two quantities are equal.

| | Column A | Column B |
|-------------|--|--|
| 65 . | The number of possible rational zeros of $f(x) = x^4 - 3x^2 + 5x + 12$ | The number of possible rational zeros of $f(x) = x^2 - 13x + 20$ |
| 66. | The greatest real zero of $f(x) = x^3 + 2x^2 - 5x - 6$ | The greatest real zero of $f(x) = x^4 + 3x^3 - 2x^2 - 6x + 4$ |

★ Challenge


Find the real zeros of the function. Then match each function with its graph.

67.
$$f(x) = x^3 + 2x^2 - x - 2$$


68.
$$g(x) = x^3 - 3x + 2$$

69.
$$h(x) = x^3 - x^2 + 2$$


Α.



В



C.



EXTRA CHALLENGE

www.mcdougallittell.com

70. CRITICAL THINKING Is it possible for a cubic function to have more than three real zeros? Is it possible for a cubic function to have no real zeros? Explain.

SOLVING QUADRATIC EQUATIONS Solve the equation. (Review 5.2 for 6.7)

71.
$$x^2 - 6x + 9 = 0$$

72.
$$x^2 + 12 = 10x - 13$$

73.
$$x - 1 = x^2 - x$$

74.
$$x^2 + 18 = 12x - x^2$$

75.
$$2x^2 - 20x = x^2 - 100$$

76.
$$x^2 - 12x + 49 = 6x - 32$$

WRITING QUADRATIC FUNCTIONS Write a quadratic function in intercept form whose graph has the given x-intercepts and passes through the given point. (Review 5.8 for 6.7)

77. *x*-intercepts:
$$-3$$
, 3 point: $(0, 5)$

78. *x*-intercepts:
$$-5$$
, 1 point: $(-2, -6)$

79. *x*-intercepts:
$$-1$$
, 5 point: $(0, 10)$

81. *x*-intercepts:
$$-12$$
, -6 point: $(9, -5)$

82. *x*-intercepts: 2, 8 point:
$$(3, -4)$$

86. SPICTURE FRAMES You have a picture that you want to frame, but first you have to put a mat around it. The picture is 12 inches by 16 inches. The area of the mat is 204 square inches. If the mat extends beyond the picture the same amount in each direction, what will the final dimensions of the picture and mat be? (Review 5.2)

Quiz 2

Self-Test for Lessons 6.4-6.6

Factor the polynomial. (Lesson 6.4)

1.
$$5x^3 + 135$$

2.
$$6x^3 + 12x^2 + 12x + 24$$

3.
$$4x^5 - 16x$$

4.
$$3x^3 - x^2 - 15x + 5$$

Find the real-number solutions of the equation. (Lesson 6.4)

5.
$$7x^4 = 252x^2$$

6.
$$16x^6 = 54x^3$$

7.
$$6x^5 - 18x^4 + 12x^3 = 36x^2$$

8.
$$2x^3 + 5x^2 = 8x + 20$$

Divide. Use synthetic division when possible. (Lesson 6.5)

9.
$$(x^2 + 7x - 44) \div (x - 4)$$

10.
$$(3x^2 - 8x + 20) \div (3x + 2)$$

11.
$$(4x^3 - 7x^2 - x + 10) \div (x^2 - 3)$$

11.
$$(4x^3 - 7x^2 - x + 10) \div (x^2 - 3)$$
 12. $(12x^4 + 5x^3 + 3x^2 - 5) \div (x + 1)$

13.
$$(x^4 + 2x^2 + 3x + 6) \div (x^3 - 3)$$

14.
$$(5x^4 + 2x^3 - x - 5) \div (x + 5)$$

Find all the real zeros of the function. (Lesson 6.6)

15.
$$f(x) = x^3 - 4x^2 - 7x + 28$$

16.
$$f(x) = x^3 - 6x^2 + 21x - 26$$

17.
$$f(x) = 2x^3 + 15x^2 + 22x - 15$$

18.
$$f(x) = 2x^3 + 7x^2 - 28x + 12$$

19. S DESIGNING A PATIO You are a landscape artist designing a patio. The square patio floor is to be made from 128 cubic feet of concrete. The thickness of the floor is 15.5 feet less than each side length of the patio. What are the dimensions of the patio floor? (Lesson 6.6)